

The study about the revised $R_{23} - P$ method for metallicity estimates based on $\sim 20,000$ SDSS galaxies

S. Y. Yin^{1,2}, Y. C. Liang¹, B. Zhang^{1,2}

¹*National Astronomical Observatories, Chinese Academy of Sciences,
 20A Datun Road, Chaoyang District, Beijing 100012, China*

²*Department of Physics, Hebei Normal University, Shijiazhuang 050016,
 China; email: syyin@bao.ac.cn, ycliang@bao.ac.cn*

Abstract. We select ~ 20000 metal-rich star-forming galaxies from the SDSS-DR2 to study the $R_{23} - P$ method suggested by Pilyugin et al. for metallicity estimates. The oxygen abundances derived from their revised $R_{23} - P$ method are ~ 0.19 dex lower than those derived from the previous one, and ~ 0.60 dex lower than the Bayesian abundances obtained by the MPA/JHU group. These abundance discrepancies strongly correlate with the P parameter and weakly depend on the $\log(R_{23})$ parameter.

1. Introduction

The chemical property of interstellar gas is important to trace the chemical evolutionary status of galaxy. Accurate abundance measurements for the ionized gas in galaxies require the determination of the electron temperature (T_e), which is usually able to be measured in metal-poor environments since the needed auroral line [O III] $\lambda 4363$ can only be detected there. The common method to estimate the oxygen abundances of metal-rich galaxies ($12 + \log(\text{O}/\text{H}) \geq 8.5$) is using empirical strong-line ratios, such as R_{23} , $R_{23} - P$, [N II]/H α etc. (Kobulnicky et al. 1999; Liang et al. 2006; Yin et al. 2006, Y06). At present, The $R_{23} - P$ method (Pilyugin et al. 2001, 2005, as P01, P05) is widely used since the derived (O/H) abundances are consistent with those derived from T_e (P05; Y06). This is specially interesting for metal-rich galaxies. However, the derived abundances from P01 and P05 show some discrepancy, which has been shown by about hundred H II regions by P05. Here we select a much larger sample of $\sim 20,000$ metal-rich star-forming galaxies from the SDSS-DR2 database to study this effect in more details. Moreover, we specially compare the abundances derived from the revised $R_{23} - P$ method given by P05 with the Bayesian abundances obtained by Tremonti et al. (2004) for these SDSS galaxies. The selected sample galaxies have $12 + \log(\text{O}/\text{H})_{\text{Bay}} > 8.5$ and $12 + \log(\text{O}/\text{H})_{\text{P05}} > 8.25$, which guarantee they are in the upper branch of metallicity.

2. Comparisons between the abundances from different methods

Previously, P01 gave the $R_{23} - P$ calibration formula for metal-rich galaxies on the basis of ~ 40 H II regions. P05 revised this calibration by using an ex-




Figure 1. Comparisons between the oxygen abundances derived from P01, P05 and Bayesian technique. **(a)** The differences between the $\log(\text{O}/\text{H})$ abundances derived from the previous (P01) and revised (P05) $R_{23} - P$ methods as a function of the oxygen abundances derived from the revised $R_{23} - P$. The differences reach ~ 0.19 dex. **(b)** The differences between the $\log(\text{O}/\text{H})$ derived from P01 and P05 as a function of $\log(R_{23})$. It is constant (~ 0.2 dex) as $\log(R_{23}) < 0.5$, but becomes smaller following the increase of the $\log(R_{23})$, which may be due to the turn-over region of the R_{23} parameter. **(c)** The abundance differences as a function of P parameter. It shows a tight correlation with P , specially for the sample with smaller value of P ($P < 0.5$), but becomes constant as $P > 0.5$. **(d), (e), (f)** show the oxygen abundance differences between the Bayesian abundances and those derived from the revised $R_{23} - P$ method of P05 as a function of $\log(\text{O}/\text{H})_{\text{P05}}$, the $\log(R_{23})$, and P parameters, respectively. **Fig.(d)** shows the ~ 0.60 dex difference. **Fig.(e)** shows similar results as **Fig.(b)**. **Fig.(f)** shows that the oxygen abundance differences have a steep linear correlation with the P parameter.

tended sample, about 104 H II regions. P05 has shown the difference between these two calibrations. Here we use a much larger sample to study this discrepancy. Figs.1a-c show the large discrepancy between the two derived abundances from P01's and P05's calibrations, and the relations of this discrepancy with $\log(R_{23})$ and P . Then, we also study the large discrepancies between the Bayesian abundances and those derived from P05's calibration, and their relations with $\log(R_{23})$ and P (Figs.1d-f). See Caption of Fig.1 for more details.

In summary, different metallicity calibrations can result in very different abundances, and we should be careful when compare the abundance estimates derived from different methods.

Acknowledgments. This work was supported by the National Science Foundation of China (NSFC) Foundation under No.10403006, 10373005, 10433010, 10573022.

References

- Kobulnicky, H.A., Kennicutt, R.C.Jr., & Pizagno, J.L., 1999, ApJ, 514,544
 Liang, Y.C., Yin, S.Y., Hammer, F. et al. 2006, ApJ, 652, 257
 Pilyugin, L.S., 2001, A&A, 369, 594 (P01)
 Pilyugin, L.S., & Thuan, T.X., 2005, ApJ, 631, 231 (P05)
 Tremonti, C.A., Heckman, T.M., Kauffmann, G., et al. 2004, ApJ, 613, 898
 Yin, S.Y., Liang, Y.C. et al., 2006, A&A (in press), astro-ph/0610068 (Y06)